## FOOD/INSECT EMERGENCY ANAPHYLAXIS CARE PLAN and MEDICATION AUTHORIZATION

Connecticut State Law and Regulations 10-212(a) require a written medication order of an authorized prescriber, (physician, dentist, optometrist, advanced practice registered nurse or physician's assistant, and for interscholastic and intramural sports only, a podiatrist) and parent/guardian written authorization, for the nurse, or in the absence of the nurse, a qualified school personnel to administer medication.

School: District/Town: ACES						
	Student Name			DOB:		
STUDENT INFORMATION	Home/Cell Phone			Grade		
	KNOWN LIFE-THREATENING ALLERGIES:PEANUTS TREE_ NUTS			History of Asthma? No Yes		
FOR	☐ MILK ☐ SOY ☐ WHEAT ☐ SHELLFISH ☐ FISH (	OTHER)	-	(Increases risk of severe reaction)		
I Z ⊢	BEE STINGS LATEX EGGS:	DTHER:				
JDEN	CONFIRMED WITH ALLERGY TESTING YES NO			Give epinephrine upon exposure		
STL	KNOWN ORAL ALLERGY SYNDROME: No Yes (list):			(before the onset of any symptoms)		
	Provide separate medication authorization if treatment indicated			If Yes		
	AFTER EXPOSURE TO KNOWN OR SUSPECTED ALLERGY & ANY OF THESE SYMPTOMS:  FOLI			FOLLOW THIS PROTOCOL:		
TREATMENT PLAN	ARWAY: Difficulty breathing, swallowing, chest tightness, wheeze THROAT: Tight, hoarse, swollen tongue, difficulty swallowing/drooling CARDIAC: Dizzy, faint, confused, pale or blue, hypotension, weak pulse &/OR  ANY COMBINATION OF SYMPTOMS FROM DIFFERENT BODY AREAS:  > Swollen lips, repetitive cough, sneezing, profuse runny nose  > Hives, itching (anywhere), swelling (e.g., eyes)  > Nausea, Vomiting, diarrhea, crampy pain  1. INJECT EPINEPHRINE IMMEDIATEL 2. Call 911 3. Lie down if able, avoid rapid upright positioning & continue monitoring 4. Give Bronchodilator/Albuterol if has asthma 5. Notify Parent/Guardian 6. Notify Prescribing Provider / PCP 7. When indicated, assist student to rise very slowly.					
EPINEPHRINE	☐ Epinephrine Auto-injector, Jr (0.15mg) IM side of thigh ☐ Epinephrine Auto-injector (0.3mg) IM side of thigh  ➤ A second dose of epinephrine can be given 5 minutes or more if symptoms persist or recur.					
EPH	Relevant Side Effects  Tachycardia Other: Medication Allergies NKDA Other:					
FPI	Medication shall be 2023-2024	NOTE: IF NURSE IS NOT AVAILABLE, THE EPINEPHRINE AUTO				
_	administered during school year:	INJECTOR MAY BE GIVEN BY DESIGNATED SCHOOL PERSONNEL WITH				
	EXPOSURE OR FOR ANY ANAPHYLAXI			YLAXIS SYMPTOMS		
T(	TO BE COMPLETED BY PARENT AND AUTHORIZED HEALTHCARE PROVIDER: REQUIRED					
	Prescriber's Authorization to Self- Administer No *Yes, Confirms student is capable PRESCRIBER'S PRINTED NAME OR ST.					
-	to safely and properly administer medication			on		
/TION	Prescriber's Signature:	Date:				
AUTHORIZATION	Parent/Guardian Consent ☐ I authorize the student to possess and self-administer medication OR ☐ I authorize this medication to be administered by school personnel  > I also authorize communication between the prescribing health care provider and school nurse necessary for allergy management and administration of this medication					
	Signature:					
		Date:				

\*TURN OVER FORM FOR INSTRUCTIONS ON ADMINISTERING EPINEPHRINE\*

## EMERGENCY ALLERGY CARE PLAN FOR STUDENT To be completed by School Nurse \_GRADE/SCHOOL: \_\_\_\_\_\_ NAME: **ALLERGIES:** GIVE EPINEPHRINE UPON EXPOSURE TO ABOVE ALLERGY OR GIVE EPINEPHRINE AT THE ONSET OF ANY OF THE BELOW SYMPTOMS IF ALLERGEN LIKELY EATEN (OR STUDENT STUNG) **SYMPTOMS OF ANAPHYLAXIS:** Insert Picture if available • Chest tightness, shortness of breath, cough, wheezing, profuse runny nose • Dizzy, faint, pale, blue, confused • Tightness and/or itching in throat, difficulty swallowing, hoarseness, drooling • Swelling of lips, tongue, throat • Itchy mouth, itchy skin, hives • Hives, itching (anywhere), swelling (e.g. face, eyes) • Nausea, vomiting, diarrhea, crampy pain EPINEPHRINE ADMINISTRATION PROTOCOL: 1. Administer Epinephrine Auto-Injector: circle one: (0.15mg 0.3mg) 2. Have someone call 911 for ambulance, don't hang up, and stay with student 3. Administer albuterol if authorized (has asthma) 4. Lie down if able; avoid rapid rise to upright position 5. Notify school and parent/quardian as soon as possible **EPI AUTO-INJECTOR DIRECTIONS:** For EPIPEN and EPIPEN JR.: 1. Stay Calm 2. Grip in your dominant hand as shown 3. Pull off blue activation cap. 4. Hold orange tip near outer thigh, OK to inject through clothing, but make sure pocket on that leg is empty. 5. Swing and jab firmly into outer thigh until you hear it click so you know it's injecting the medicine. Hold in place and count to 10; remove and massage 10 sec. (orange tip will automatically slide over needle)

EMERGENCY CONTACTS	EMERGENCY/PHYSICIAN CONTACTS
Name:	Name:
Relation:	Phone:
Phone:	

6. Auto-injector should be given to EMS to take to E.R.