#### FOOD/INSECT EMERGENCY ANAPHYLAXIS CARE PLAN and MEDICATION AUTHORIZATION

Connecticut State Law and Regulations 10-212(a) require a written medication order of an authorized prescriber, (physician, dentist, optometrist, advanced practice registered nurse or physician's assistant, and for interscholastic and intramural sports only, a podiatrist) and parent/guardian written authorization, for the nurse, or in the absence of the nurse, a qualified school personnel to administer medication.

Sch	ool: Distric	ct/Town:	ACES	
	Student Name			DOB:
NO	Home/Cell Phone			Grade
STUDENT INFORMATION			\$	History of Asthma? No Yes
INFO	☐ MILK ☐SOY ☐ WHEAT ☐SHELLFISH ☐FISH (( ☐ BEE STINGS ☐LATEX ☐ EGGS:	•		
DENT				Give epinephrine upon exposure
STUI	KNOWN ORAL ALLERGY SYNDROME: ON OYes (list	t);		(before the onset of any symptoms)
	<ul> <li>Provide separate medication authorization if tr</li> </ul>		ed	If Yes
	· · · · · · · · · · · · · · · · · · ·		<del></del>	
	AFTER EXPOSURE TO KNOWN OR SUSPECT & ANY OF THESE SYMPTOMS:	<u>[ED ALLERGY</u> :		FOLLOW THIS PROTOCOL:
TREATMENT PLAN	AIRWAY: Difficulty breathing, swallowing, chest tightnes THROAT: Tight, hoarse, swollen tongue, difficulty swallo CARDIAC: Dizzy, faint, confused, pale or blue, hypotens &/OR	owing/drooling		<ol> <li>INJECT EPINEPHRINE IMMEDIATELY!</li> <li>Call 911</li> <li>Lie down if able, avoid rapid upright positioning &amp; continue monitoring</li> </ol>
ATM	ANY COMBINATION OF SYMPTOMS FROM DIFFERE		<u>; TV '</u>	4. Give Bronchodilator/Albuterol if has asthma
TRE,	<ul> <li>Swollen lips, repetitive cough, sneezing, profuse run</li> <li>Hives, itching (anywhere), swelling (e.g., eyes)</li> </ul>	ny nose	!	5. Notify Parent/Guardian
	<ul> <li>Nausea, Vomiting, diarrhea, crampy pain</li> </ul>			<ol> <li>Notify Prescribing Provider / PCP</li> <li>When indicated, assist student to rise very slowly.</li> </ol>
			·	
	Epinephrine Auto-injector, Jr (0.15mg) IM side of thig	Jh 🗌 Epinephr	ine Auto-inj	ector (0.3mg) IM side of thigh
RINE	> A second dose of epinephrine can be given 5	minutes or more	if symptor	ns persist or recur.
EPINEPHRINE	Relevant Side Effects 🗌 Tachycardia 📋 Other:	Medicat	tion Allergi	es 🗌 NKDA 📋 Other:
Medication shall be administered during school year: 2022 2023 NOTE: IF NURSE IS NOT AVAILABLE, THE EPINEPH INJECTOR MAY BE GIVEN BY DESIGNATED SCHOOL				•

#### TO BE COMPLETED BY PARENT AND AUTHORIZED HEALTHCARE PROVIDER: REQUIRED

EXPOSURE OR FOR ANY ANAPHYLAXIS SYMPTOMS

	Prescriber's Authorization to Self- Adr	ninister No *Yes, Confirms student is capable to safely and properly administer medication	PRESCRIBER'S PRINTED NAME OR STAMP
TION	Prescriber's Signature:	Date:	
AUTHORIZA	🗌 I authorize	the student to <b>possess</b> and <b>self-administer</b> medication <b>OR</b> this medication to be <b>administered by school personnel</b> en the prescribing health care provider and school nurse dministration of this medication	
	Signature:	Date:	

#### \*TURN OVER FORM FOR INSTRUCTIONS ON ADMINISTERING EPINEPHRINE\*

## EMERGENCY ALLERGY CARE PLAN FOR STUDENT To be completed by School Nurse

GRADE/SCHOOL: \_\_\_\_\_

#### ALLERGIES:

NAME:

#### GIVE EPINEPHRINE UPON EXPOSURE TO ABOVE ALLERGY OR

**GIVE EPINEPHRINE AT THE ONSET OF ANY OF THE BELOW SYMPTOMS IF ALLERGEN LIKELY EATEN (OR STUDENT** STUNG)

## SYMPTOMS OF ANAPHYLAXIS:

- Chest tightness, shortness of breath, cough, wheezing, profuse runny nose
- Dizzy, faint, pale, blue, confused
- Tightness and/or itching in throat, difficulty swallowing, hoarseness, drooling
- Swelling of lips, tongue, throat
- Itchy mouth, itchy skin, hives
- Hives, itching (anywhere), swelling (e.g. face, eyes)
- Nausea, vomiting, diarrhea, crampy pain

# EPINEPHRINE ADMINISTRATION PROTOCOL:

- 1. Administer Epinephrine Auto-Injector: circle one: (0.15mg 0.3mg)
- 2. Have someone call 911 for ambulance, don't hang up, and stay with student
- 3. Administer albuterol if authorized (has asthma)
- 4. Lie down if able; avoid rapid rise to upright position
- 5. Notify school and parent/guardian as soon as possible

# EPI AUTO-INJECTOR DIRECTIONS:

## For EPIPEN and EPIPEN JR.:

- 1. Stay Calm
- 2. Grip in your dominant hand as shown
- 3. Pull off blue activation cap.
- 4. Hold orange tip near outer thigh, OK to inject through clothing, but make sure pocket on that leg is empty.
- 5. Swing and jab firmly into outer thigh until you hear it click so you know it's injecting the medicine. Hold in place and count to 10; remove and massage 10 sec. (orange tip will automatically slide over needle)
- 6. Auto-injector should be given to EMS to take to E.R.

## **EMERGENCY CONTACTS**

## **EMERGENCY/PHYSICIAN CONTACTS**

Name:

Relation:

Phone:

Name:

Phone:



	Not		
2	×		and the second se
	J	-	2

Insert Picture if

available